

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 249 237 A1

(12)

## EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:  
16.10.2002 Bulletin 2002/42

(51) Int Cl.7: **A61K 31/22**, A61K 7/00,  
A61P 17/00, A61P 43/00

(21) Application number: **00981789.1**

(86) International application number:  
**PCT/JP00/08942**

(22) Date of filing: **15.12.2000**

(87) International publication number:  
**WO 01/045697 (28.06.2001 Gazette 2001/26)**

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE TR**  
Designated Extension States:  
**AL LT LV MK RO SI**

(30) Priority: **21.12.1999 JP 36237799**  
**30.03.2000 JP 2000093152**  
**24.10.2000 JP 2000323637**  
**09.11.2000 JP 2000341577**

(71) Applicant: **Kanebo Limited**  
**Sumida-ku, Tokyo 131-0031 (JP)**


(72) Inventors:  
• **UCHIWA, Hideyo**  
**Odawara-shi, Kanagawa 250-0055 (JP)**  
• **HIROTSU, Sachiyo**  
**Odawara-shi, Kanagawa 250-0002 (JP)**

- **HARATAKE, Akinori**  
**Odawara-shi, Kanagawa 250-0878 (JP)**
- **IKEMOTO, Takeshi**  
**Minamiasigara-shi, Kanagawa 250-0013 (JP)**
- **MATSUI, Junichi**  
**Odawara-shi, Kanagawa 250-0005 (JP)**
- **YAMAZAKI, Shunsuke**  
**Odawara-shi, Kanagawa 250-0877 (JP)**
- **YOSHIDA, Masaki**  
**Odawara-shi, Kanagawa 250-0816 (JP)**
- **HAYASE, Motoi**  
**Odawara-shi, Kanagawa 250-0055 (JP)**

(74) Representative: **Prins, Adrianus Willem et al**  
**Vereenigde,**  
**Nieuwe Parklaan 97**  
**2587 BN Den Haag (NL)**

## (54) SKIN EXTERNAL AGENTS AND DRUGS

(57) The present invention provides an epidermal permeability barrier function-enhancing agent, an amphiregulin expression inhibitor, an epidermal hyperproliferation inhibitor, a heparin-binding epidermal growth factor-like growth factor (HB-EGF) expression inducing agent, an ameliorating agent for aged skin, an agent to be used together with retinoic acid or a derivative thereof, an agent to prevent dry skin caused by retinoic acid, and a skin external agent composition, which all comprises fumaric acid diester derivatives expressed by the following general formula (1)



wherein R<sup>1</sup> represents a C<sub>1-4</sub> alkylene group; R<sup>2</sup> and R<sup>3</sup> each represents a linear or branched C<sub>1-8</sub> alkyl group; and n is an integer of 2 to 5.

**Description**

## Field of the Invention

- 5 **[0001]** The present invention relates to skin external agents such as cosmetics and medicines. Particularly, the present invention relates to an epidermal permeability barrier function-enhancing agent which enhances the epidermal permeability barrier function against environmental influences such as sunburn and quickly ameliorates damaged function of epidermal permeability barrier such as dry skin to maintain the skin in a dermatological and cosmetologically healthy state.
- 10 **[0002]** The present invention relates also to an amphiregulin expression inhibitor and an epidermal hyperproliferation inhibitor which are useful for therapy, prevention, and studies on various diseases involving excessive amphiregulin expression and dry skin.
- [0003]** Further, the present invention relates to an ameliorating agent for aged skin which prevents metabolic disruption and thinning of the skin due to aging and is effective in treatment of age spots. Still further, the present invention
- 15 relates to an agent that is used together with retinoic acid or a derivative thereof in order to prevent side effects, such as dry skin, caused by retinoic acid or a derivative thereof.

## Description of the Prior Art

- 20 **[0004]** All organisms including the human being are influenced by their external environment. However, higher animals such as mammal have an organ which minimizes the environmental influence on other important organs and systems for life. The organ is the skin that has, as its most important function, an epidermal permeability barrier function to prevent foreign substances from invading into the body and excessive vaporization of moisture from the interior of the body.
- 25 **[0005]** The epidermal permeability barrier function can temporarily disrupt due to diseases or exposure to organic solvents, surfactants or UV-rays, which disturbs the skin's internal environments. If this condition continues, such a risk may arise that pathological bacteria and harmful chemical substances may invade through the skin into the interior of the body. Accordingly, it is necessary to quickly restore the epidermal permeability barrier function. In addition, in the state of disruption in the epidermal permeability barrier function, the epidermis becomes dry and its surface is covered with scales, which leads to a cosmetologically undesirable look.
- 30 **[0006]** The epidermal permeability barrier function is formed and maintained by keratinocyte turnover and keratinization near the stratum corneum. Where the epidermal permeability barrier function disrupts, hyperproliferation of keratinocytes occurs and continuous hyperproliferation is considered to diminish normal keratinocyte turnover and keratinization, which delays the recovery of the barrier function.
- 35 **[0007]** Psoriasis is a disease with unknown causes, and is accompanied with hyperproliferation of keratinocytes and keratinization insufficiency. Drugs such as steroids and Cyclosporin A, immunosuppressive agent, are generally administered for its treatment. However, it is pointed out that these drugs are dangerous for routine or daily application. Therefore, a drug is desired which reduces and normalizes epidermal hyperproliferation more safely. Similarly, a safe and effective external agent is in demand for dermatological diseases such as atopic dermatitis accompanied with
- 40 disruption of the epidermal permeability barrier function.
- [0008]** Appearance of the skin is largely influenced by the skin's overall metabolic ability and its conditions. For example, when the epidermal permeability barrier function is disrupted, the skin surface becomes dry and is covered with scales, which is an undesirable state. By hypometabolism with aging, change in skin tone caused by thinning of the epidermis, dullness, age spot, and wrinkles, lead to loss of beauty in appearance.
- 45 **[0009]** The appearance of a beautiful skin is maintained by active keratinocyte turnover and normal keratinization near the stratum corneum. When the keratinocyte turnover is lowered, keratinization may not proceed smoothly, resulting in dry skin. Lowered turnover due to aging results in dullness in the skin or age spots caused by accumulated melanin. The epidermal turnover is considered to depend on a cellular mitosis rate in the basal layer, and a decreased mitosis rate due to aging results in thinner epidermis and wrinkles as well.
- 50 **[0010]** The keratinocyte turnover is considered to be controlled by the expression of epidermal growth factors such as amphiregulin, heparin-binding EGF-like growth factor (HB-EGF), and TGF- $\alpha$ . In particular, HB-EGF expressing near the basal layer is considered to be the greatest contributing factor and, therefore, the epidermal turnover is up-regulated by inducing the HB-EGF expression [Journal of Investigative Dermatology (1998), Vol. 111, pp. 715 to 721].
- [0011]** Retinoic acid is known as a very effective external agent and is also known to induce HB-EGF expression.
- 55 On the other hand, retinoic acid has side effects such as dry skin and temporary darkening, so that, if it is used as a skin topical agent, physicians must monitor the patients very carefully. Therefore, patients who need routine prevention and treatments desire the development of safer drugs and a means to prevent such side effects.
- [0012]** It is known that the expression of amphiregulin does not occur in the skin showing normal turnover, but is

restricted in the epidermis exhibiting hyperproliferation or proliferation in cultured keratinocytes. Consequently, it is believed that the epidermal hyperproliferation can be inhibited specifically and effectively and, accordingly, functions of the skin are normalized by inhibiting the expression of amphiregulin. The expression of amphiregulin is involved not only in the hyperproliferation of the skin, but also in the process of development of skin cancer according to a report. However, any substance is not yet known which specifically inhibits the expression of amphiregulin. Therefore, also for the purpose of research progress, the development of a drug which inhibits the expression of amphiregulin is strongly desired.


**[0013]** The inventors found that dimethyl fumarate and diethyl fumarate are effective for ameliorating the epidermal permeability barrier function (Japanese Patent Application Laid-Open No. 12-178116/2000). In addition, it is known that fumaric acid and fumaric acid esters have a sebum secretion-promoting function (Japanese Patent Application Laid-Open No. 4-18017/1992), and fumaric esters are orally administrated for the treatment of psoriasis (British Journal of Dermatology 1998:138:456-460, 1999:141:424-429). However, it is also known that, when they are used externally, perilesional skin irritation, macular papular rashes, and urticarial reactions are caused and there is a risk of causing sensitization and contact urticaria (Dermatology 1994:188:126-130).

#### Disclosure of the Invention

**[0014]** The purpose of the present invention is to provide an epidermal barrier function-enhancing agent and a composition for skin external application which are very safe, quickly ameliorate the epidermal permeability barrier function disruption, and have excellent effects on maintaining the skin in a cosmetologically healthy state; a HB-EGF expression-inducing agent to accelerate epidermal turnover; an amphiregulin expression inhibitor to diminish epidermal hyperproliferation; an agent to be used together with retinoic acid or a derivative thereof to prevent dry skin caused by retinoic acid or a derivative thereof; and an ameliorating agent for aged skin which ameliorates hypometabolism and thinning of the skin due to aging.

**[0015]** To attain the aforementioned purposes, the inventors earnestly have investigated effects and safety on skin of various kinds of fumaric acid diester derivatives, and have found that the fumaric acid diester derivatives expressed by the following general formula (1) are safe to skin (i.e., not showing sensitization) and have an excellent epidermal permeability barrier-function enhancing function. The inventors have also found that a skin external agent composition containing this derivative has excellent effects of ameliorating the epidermal permeability barrier function; maintaining the skin in a dermatologically and cosmetologically healthy state; inducing the expression of HB-EGF; inhibiting the expression of amphiregulin; preventing skin thinning due to aging; and preventing dry skin caused by retinoic acid.

**[0016]** The present invention provides an epidermal permeability barrier function-enhancing agent, an amphiregulin expression inhibitor, an epidermal hyperproliferation inhibitor, a heparin-binding epidermal growth factor-like growth factor (HB-EGF) expression inducing agent, an ameliorating agent for aged skin, an agent to be used together with retinoic acid or a derivative thereof, an agent to prevent dry skin caused by retinoic acid, and a skin external agent composition, which all comprises fumaric acid diester derivatives expressed by the following general formula (1)



wherein  $\text{R}^1$  represents a  $\text{C}_{1-4}$  alkylene group;  $\text{R}^2$  and  $\text{R}^3$  each represents a linear or branched  $\text{C}_{1-8}$  alkyl group; and  $n$  is an integer of 2 to 5.

The present invention also provides a skin external agent composition comprising the aforesaid fumaric acid diester derivatives, and retinoic acid or a derivative thereof.

#### Description of the Preferred Embodiments

**[0017]** The embodiments of the present invention will be described below in detail.

**[0018]** Examples of the fumaric acid diester derivatives expressed by the aforementioned general formula (1) used in the present invention include bis (diethylene glycol monoethyl ether) fumarate expressed by the following general



**[0029]** In addition to the substances mentioned above, vehicles, surfactants and antioxidants allowed to be used in medicines, quasi-drugs or cosmetics can be blended in the present drugs or the skin external agent compositions in such an amount that the purpose of the present invention is achieved.

## 5 Examples

**[0030]** The present invention will be explained with reference to the following Tests, Examples and Comparative Examples. Test 1 (Epidermal Permeability barrier Function Amelioration Test)

**[0031]** An effect of improving epidermal permeability barrier function by applying the agents on the skin of which epidermal permeability barrier function has been disrupted was evaluated by the following test method.

### 1. Experimental animals used

**[0032]** Male hairless mice (Hos: Hr-1) of 10 weeks old at the beginning of the test were used with each 5 mice per group.

### 2. Measurement of the epidermal permeability barrier state

#### 2-1. Measurement device and conditions

**[0033]** Transepidermal water loss (TEWL) was measured as follows using a continuous perspiration measurement device, Hydrograph AMU-100 (ex K & S Co., Ltd.). A 1cm<sup>2</sup> capsule was closely attached to the skin and nitrogen gas was led into the capsule in a rate of 300mL/min. The amount of water vapor in the nitrogen gas was measured before the gas was fed to the capsule and after it was recovered from the capsule. From a difference between the amounts, the amount of water in mg\_vaporized from 1cm<sup>2</sup> area of the skin per minute, TEWL, was calculated.

#### 2-2. Samples and the experimental method

**[0034]** The samples were prepared by mixing the compounds indicated in Table 1 below, e.g., bis(diethylene glycol monoethyl ether) fumarate, with a 50% aqueous ethanol solution (base). The TEWL of the hairless mouse dorsal skin was first measured at the beginning of the test. Subsequently, 0.05mL of each sample was applied on an area of approximately 2.5cm diameter in the dorsal skin of the hairless mice once a day from Monday to Friday (pre-application). Then, on the third day from the last day of the pre-application, the skin was irradiated once with ultraviolet B ray (UVB) at a dose of 0.15J/cm<sup>2</sup>. Each sample was applied immediately after the UVB irradiation until the 3<sup>rd</sup> day from the irradiation in the same amount and at the same frequency as in the pre-application. The TEWL was measured on the 3<sup>rd</sup> and 4<sup>th</sup> days after the irradiation. TEWL change rates (= TEWL value on the 3<sup>rd</sup> or 4<sup>th</sup> day after the irradiation/TEWL value at the beginning of the test) were calculated, which show changes in TEWL relative to the initial TEWL due to the exposure to UVB, and the mean values thereof were compared between the control group and each sample group.

#### 2-3. A method of assessment of the effects

**[0035]** TEWL changes with the degree of disruption of the epidermal permeability barrier function. A higher value of TEWL change rate implies a greater degree of disruption of the epidermal permeability barrier function. In the case of the application of the base alone, disruption in the epidermal permeability barrier function by UVB was similar to that in the case of non-application. Therefore, a group showing a lower TEWL change rate than that of the control group, where the base alone was applied and thus showed the natural state of the disruption in the epidermal permeability barrier function, was assessed as available in the epidermal permeability barrier function amelioration.

**[0036]** The TEWL change rates in the hairless mice after the application of the samples are as shown in Table 1.

(Table 1)

Compounds	Concentrations, %	TEWL change rate	
		3 <sup>rd</sup> day	4 <sup>th</sup> day
Bis (diethylene glycol monoethyl ether) fumarate	1	3.25	2.49
-	0	7.98	7.27

(Table 1) (continued)

Compounds	Concentrations, %	TEWL change rate	
		3 <sup>rd</sup> day	4 <sup>th</sup> day
Diphenylethyl fumarate	1	7.82	6.57
Dimethyl fumarate	1	3.24	2.50
Mean values			

[0037] As is seen from the results of this test, bis (diethylene glycol monoethyl ether) fumarate which is a fumaric acid diester derivative according to the present invention clearly reduces disruption in the epidermal permeability barrier function by UVB and ameliorates epidermal permeability barrier function, compared to the base alone and diphenylethyl fumarate, and its epidermal permeability barrier function amelioration effects is as great as that of dimethyl fumarate which has sensitizing potential. Next, a similar test was carried out using bis (diethylene glycol monoethyl ether) fumarate in various concentrations as described in Table 2 below.

(Table 2)

Compounds	Concentrations, %	TEWL change rate	
		3 <sup>rd</sup> day	4 <sup>th</sup> day
Bis (diethylene glycol monoethyl ether) fumarate	0.001	7.01	7.31
ditto	0.01	6.46	6.84
ditto	0.1	3.79	5.21
ditto	1	3.62	4.33
-	0	8.56	8.11
Mean values			

#### Test 2 [Safety (sensitization) Test]

[0038] Safety (sensitization) was evaluated by a maximization test. An area of 4x6cm<sup>2</sup> of skin on the scapula of a Hartley strain guinea pig (female) weighing 350 to 400g was shaved and given intradermal injections with each three injections in two lines according to the following procedures:

- (1) each 0.05 mL of Freund's Complete Adjuvant (hereinafter simply called FCA) was intradermally injected at each one spot in the right and left sites.
- (2) each 0.05mL of a 3% fumaric acid diester derivative solution in ethanol of as a test sample was intradermally injected at another each one spot in the right and left sites.
- (3) to a 6% fumaric acid diester derivative solution in FCA as a test sample, the same amount of sterilized water was added and resultant emulsified solution was intradermally injected at a right side and a left side sites in an amount of 0.05mL.

[0039] At one week after the aforementioned procedures, the skin of the same sites was shaved and 10% of sodium lauryl sulfate in vaseline was applied to cause light inflammation. After 24 hours from the application, 0.2mL of a 3% fumaric diester derivative solution in ethanol as a test sample, which was applied on a piece of gauze, was patched close on the same sites for 48 hours. On the 21<sup>st</sup> day after the intradermal injection, the abdominal area was shaved, to which a solution of a fumaric acid diester derivative in ethanol as a test sample, which was applied in a piece of gauze, was patched close for 24 hours. The symptoms in the skin were visually examined according to the following criteria after 48 hours.

Symptoms	Rating
No reaction	0
Scattered mild redness	1
Moderate and diffuse redness	2

(continued)

Symptoms	Rating
intense redness and swelling	3

## 2. Test results

[0040] The sensitization test results with various fumaric acid diester derivatives after 48 hours are shown below. As seen from the test results, bis (diethylene glycol monoethyl ether) fumarate, which is a fumaric acid diester of the present invention, was confirmed to have no sensitizing potential. However, bis (ethylene glycol monomethyl ether) fumarate which has the general formula (1) with  $n=1$  was found to have sensitizing potential as diethyl fumarate did.

Test samples	Positivity
Diethyl fumarate	90%
Bis (ethylene glycol monomethyl ether) fumarate	90%
Bis (diethylene glycol monoethyl ether) fumarate	0%

## Test 3 (Human dry skin amelioration test)

[0041] A skin lotion having the composition described below was prepared by the formulation method described below. The sample lotion was applied on skin at an inside of an upper arm of 10 healthy subjects (males of 25 to 52 years old) and tested for human dry skin amelioration (epidermal permeability barrier function recovery test). As a control, one having the same composition except for bis (diethylene glycol monoethyl ether) fumarate was used.

Skin lotion composition	Amount, %
Component (A)	
Olive oil	10.0
Isopropyl myristate	1.0
Polyoxyethylene (15) nonylphenyl ether	0.5
Propylene glycol	1.0
Glycerin	2.0
Component (B)	
Methylparaben	0.1
Ethanol	7.0
Bis (diethylene glycol monoethyl ether) fumarate	1.0
Purified water	Balance

## Preparation method

[0042] Components (A) and (B) were separately dissolved uniformly, and then were combined to disperse with stirring. The mixture was filled in a container. The contents were shaken to effect uniform dispersion before each use.

[0043] Each sample was applied to the test site (2cm x2cm for each sample) in an amount of 0.1mL once a day after bathing, which was continued for 2 months (61 days). On the 62<sup>nd</sup> day from the beginning of the application, the skin surface was wiped with acetone to attain 0.15mg/cm<sup>2</sup>/min of TEWL, indicating an induced state of dry skin. Subsequently, the sample was applied again in the same way as described above. Along with investigating how much the epidermal permeability barrier function recovers, and also the skin surface status and the presence of side effects such as rashes and itching were examined.

[0044] The skin lotion with the aforementioned composition according to the present invention clearly demonstrated the dry skin amelioration effect compared to the control. Further, no skin abnormalities such as rashes and dryness were observed.

## Test Example 4 (Test of measuring an HB-EGF expression rate)

[0045] The HB-EGF expression rates were investigated by the following test methods where the compounds of the present invention were applied on the keratinocytes.

## 1. Keratinocytes used in the test

[0046] Cultured keratinocytes of human epidermis origin, commercially available from Kurabo Corporation, were used.

## 2. Measurement of HB-EGF expression in the keratinocytes

## (1) Experimental method

[0047] Keratinocyte suspension was adjusted to a concentration of  $400 \times 10^4$  units/mL on an MCDB153 culture medium supplemented with BPE (bovine pituitary extract), insulin, ethanolamine, phosphoethanolamine, and hydrocortisone as growth promoters. Four milliliters of the cell suspension was inoculated to a 60mm plate (ex Falcon Corporation). Under an atmosphere of 95%(v/v) air and 5% (v/v) carbon dioxide gas, a confluent state was achieved by static culturing at 37°C for 3 days, while refreshing the culture media every other day.

[0048] After removing the culture medium supernatant by suction, the culture medium was replaced with an MCDB 153 culture medium (containing insulin) to which dimethyl fumarate had been added at a final concentration of 50 $\mu$ mol/L or bis (diethylene glycol monoethyl ether) fumarate in 300 $\mu$ mol/L. This plate was subjected to static culturing at 37°C for 24 hours under an atmosphere consisting of 95% (v/v) air and 5% (v/v) carbon dioxide gas. Using a QIAGEN™ RNA kit, whole mRNA was extracted. Each amount of HB-EGF mRNA and G3PDH mRNA per total mRNA was determined by Northern blotting. The amount of each mRNA per total in the absence of the fumarate was measured as the control. The amount of mRNA in the presence of the fumarate was expressed in % relative to those of the control.

## (2) Results

[0049] The relative amounts of mRNA for HB-EGF and G3PDH in the presence of each sample are as shown below.

Compounds added	Amount of mRNA ,% relative to the control	
	HB-EGF	G3PDH
Bis(diethylene glycol monoethyl ether) fumarate	380%	98%
Dimethyl fumarate	470%	105%

[0050] As can be seen from the above-mentioned results, bis (diethylene glycol monoethyl ether) fumarate induced the expression of HB-EGF as strongly as dimethyl fumarate which has sensitizing potential.

## Test Example 5 (Test on capability of inhibiting reduction in epidermis thickness)

[0051] The following test method was used in order to investigate inhibition of reduction in the epidermis thickness when the compound of the present invention was applied on aged skin.

## 1. Experimental animals used in the test

[0052] Experimental animals used included 2 groups of aged hairless mice of 33 weeks old at the beginning of the test with each 5 mice per group, and one group of 5 hairless mice of 9 weeks old at the beginning of the test.

## 2. Measurement of the epidermis thickness

## (1) Samples and the experimental method

[0053] A sample was prepared by adding bis (diethylene glycol monoethyl ether) fumarate to a 50% ethanol aqueous solution (base) at a concentration of 0.5%. This sample was applied in an amount of 0.05mL on the dorsal skin of hairless mice (in a diameter of approximately 2.5cm) once a day for consecutive 5 days per week followed by 2 days



interval. This schedule was repeated for 4 weeks. Subsequently, a portion of the skin was sampled and a frozen section was prepared. The thickness of the epidermis was measured by microscopic observation. The mean values of the epidermis thicknesses in hairless mice after the application of the fumarate are shown below.

Compound (Amount added)	Mean values of the epidermis thickness, $\mu\text{m}$
(33 weeks old hairless mice)	
Bis (diethylene glycol monoethyl ether) fumarate (0.5%)	20.96
Untreated	17.24
(9 weeks old hairless mice)	
Untreated	28.88

The results of this test indicates that thinning of the epidermis due to aging was clearly inhibited by the addition of bis (diethylene glycol monoethyl ether) fumarate compared to the non-treatment.

Test Example 6 (Test of inhibition of the expression of amphiregulin)

**[0054]** The amount of expression of amphiregulin in the keratinocytes was determined by the following test method.

1. Keratinocytes used in the test

**[0055]** Cultured keratinocytes of human epidermis origin (ex Kurabo Corporation) were used.

2. Measurement of amphiregulin expression in the keratinocytes

(1) Experimental method

**[0056]** Keratinocyte suspension was adjusted to a concentration of  $150 \times 10^4$  units/mL in an MCDB153 culture medium supplemented with BPE (bovine pituitary extract), insulin, ethanolamine, phosphoethanolamine, and hydrocortisone as growth promoters. Four milliliters of the cell suspension was inoculated to a 60mm plate (ex Falcon Corporation). Under an atmosphere of 95%(v/v) air and 5% (v/v) carbon dioxide gas, a confluent state was achieved by static culturing at 37°C for 1 day.

After removing the culture medium supernatant by suction, the culture medium was replaced with an MCDB 153 culture medium (containing insulin) to which bis (diethylene glycol monoethyl ether) fumarate had been added in a final concentration of 300 $\mu\text{mol/L}$  or dimethyl fumarate in 70 $\mu\text{mol/L}$ . This plate was subjected to static culturing at 37°C for 24 hours under an atmosphere consisting of 95%(v/v) air and 5% (v/v) carbon dioxide gas. Using a QIAGEN<sup>TM</sup> RNA kit, whole mRNA was extracted. Each amount of amphiregulin and G3PDH mRNA per total mRNA was determined by Northern blotting. The amount of each mRNA per total in the absence of the fumarate was measured as the control. The amount of mRNA in the presence of the fumarate was expressed in % relative to those of the control.

(2) Results

**[0057]** The relative amounts of amphiregulin mRNA and G3PDH mRNA in the presence of each sample are as shown below.

Agents added	Amount of mRNA ,% relative to the control	
	Amphiregulin	G3PDH
Example 1. Bis (diethylene glycol monoethyl ether) fumarate	8%	102%
Comparative Example 1 Dimethyl fumarate	39%	98%

The fumaric acid diester derivative of the present invention was found to inhibit the amphiregulin expression as strongly as dimethyl fumarate having sensitizing potential does. Since there was no difference in the amount of G3PDH mRNA between the control and Example 1, the mRNA expression inhibition action by the fumarate diester derivative was

found to be specific to amphiregulin.

Test Example 7 (Test of inhibition of amphiregulin expression when used together with retinoic acid)

5 **[0058]** The amount of amphiregulin expression in the presence of retinoic acid in the keratinocytes was determined by the following method.

1. Keratinocytes used in the present test

10 **[0059]** Cultured keratinocytes of human epidermis origin (ex Kurabo Corporation) were used.

2. Measurement of amphiregulin expression in the keratinocytes

(1) Experimental method

15 **[0060]** Keratinocyte suspensions was adjusted to a concentration of  $400 \times 10^4$  units/mL in an MCDB153 culture medium supplemented with BPE (bovine pituitary extract), insulin, ethanolamine, phosphoethanolamine, and hydrocortisone as growth promoters. Four milliliters of the cell suspension was inoculated to a 60mm plate (ex Falcon Corporation). Under an atmosphere of 95%(v/v) air and 5% (v/v) carbon dioxide gas, a confluent state was achieved by static culturing at 37°C for 3 days, while refreshing the culture every other day.

20 After removing the culture medium supernatant by suction, the culture medium was replaced with an MCDB 153 culture medium (containing insulin) to which bis (diethylene glycol monoethyl ether) fumarate at a final concentration of  $300 \mu\text{mol/L}$  or dimethyl fumarate at  $70 \mu\text{mol/L}$ , and optionally retinoic acid at  $1 \mu\text{mol/L}$  had been added. This plate was subjected to static culturing at 37°C for 24 hours under an atmosphere consisting of 95%(v/v) air and 5% (v/v) carbon dioxide gas. Using a QIAGEN™ RNA kit, whole mRNA was extracted. Each amount of amphiregulin mRNA and G3PDH mRNA per total mRNA was determined by Northern blotting. The amount of each mRNA per total in the absence of the fumarate was measured as the control. The amount of mRNA in the presence of the fumarate was expressed in % relative to those of the control.

30 (2) Results

**[0061]** The relative amounts of amphiregulin mRNA and G3PDH mRNA after the addition of each sample are as shown below.

Agents added	Amount of mRNA, % relative to the control	
	Amphiregulin	G3PDH
Example 2. Bis (diethylene glycol monoethyl ether) fumarate + retinoic acid	70%	102%
40 Comparative Example 2. Dimethyl fumarate + retinoic acid	120%	98%
Comparative Example 3. Retinoic acid alone	300%	105%

45 As seen in Comparative Example 3, amphiregulin was strongly induced by the addition of retinoic acid and the induced amphiregulin is considered to be closely involved in formation of dry skin caused by retinoic acid. Example 2 reveals that the fumaric acid diester derivatives of the present invention more strongly inhibit the expression of amphiregulin induced by the addition of retinoic acid than dimethyl fumarate having sensitizing potential does. Since the amount of mRNA for G3PDH as the control in Example 2 agreed those in the Comparative Examples within an experimental error, the function of the fumaric acid diester derivative to inhibit the expression of mRNA was found to be specific to amphiregulin.

Test Example 8 (Test of inhibition of dry skin)

55 **[0062]** The effect of inhibiting dry skin was investigated by the following test method in which the present fumarate was applied to the dry skin induced by retinoic acid.

## 1. Experimental animals used in the test

[0063] Hairless mice of 10 weeks old at the beginning of the test were used with five mice per group.

## 2. Measurement of TEWL

[0064] A sample was prepared by adding dimethyl fumarate or bis (diethylene glycol monoethyl ether) fumarate to a 50% ethanol aqueous solution (base) in a concentration of 0.5%. This sample was applied in an amount of 0.05mL to the dorsal skin of hairless mice (in a diameter of approximately 2.5cm) once a day for consecutive 5 days (pre-application). Subsequently, 0.05mL of a 0.05% retinoic acid solution in 50% ethanol aqueous solution was applied once a day for 4 days. During these 4 days, retinoic acid was applied in the morning, and, some hours later in the afternoon, the sample was applied. The TEWL values were measured by a Hydrograph AMU-100 (ex K & S Corp.) after completion of the application. In Comparative Example 5, a control sample comprising the base alone was used instead of the sample. In Comparative Example 6, the base was always applied. The TEWL values after the application of the samples are as shown below.

Group	Components	Mean value of TEWL , mg/cm <sup>2</sup> /min.
Example 3	Bis (diethylene glycol monoethyl ether) fumarate	0.055
Comparative Example 4	Dimethyl fumarate	0.063
Comparative Example 5	Retinoic acid, and no fumarate	0.110
Comparative Example 6	Always the base only	0.012

[0065] As seen above, increase in TEWL due to the application of retinoic acid was observed in Comparative Example 5 compared with Comparative Example 6 where the base alone was applied. In Example 3, because of the addition of the fumaric acid diester derivative of the present invention, apparently smaller increase in TEWL than in Comparative Example 5 was observed, which indicates amelioration of dry skin induced by retinoic acid. The TEWL value in Example 3 was almost equal to that in Comparative Example 4 where dimethyl fumarate having sensitizing potential was used.

Test Example 9 (Test of depression of UV-induced epidermal hyperproliferation)

[0066] The inhibition of epidermal hyperproliferation was investigated by the following test method in which the present fumarate was applied to the skin where epidermal hyperproliferation had been induced by UV irradiation

## 1. Experimental animals used in this test example

[0067] Hairless mice of 10 weeks old at the beginning of the test were used with 5 mice per group.

## 2. Measurement of epidermis thickness

## (1) Samples and experimental method

[0068] Samples were prepared by adding dimethyl fumarate or bis (diethylene glycol monoethyl ether) fumarate to a 50% ethanol aqueous solution (base) at a concentration of 0.5%. Initially, this sample was applied in an amount of 0.05mL to the dorsal skin of the hairless mice (in a diameter of approximately 2.5cm) once a day and 5 times per week for 4 weeks (pre-application). Subsequently on the 3<sup>rd</sup> day from the last application of the pre-application, the skin was irradiated with a UVB ray once at a dose of 0.15J/cm<sup>2</sup>. The animals were slaughtered on the 4<sup>th</sup> day after the irradiation and skin section were made. From the section, tissue section samples were prepared by HE staining. The samples prepared were photographed using a light microscope. In the microscopic photographs, the thickness of the epidermis was measured. Using the epidermis thickness at the beginning of the test as a standard, the mean values were compared between the control group and the treated groups.

[0069] The mean values of thickness of the epidermis after the application of each sample are as shown below.

Agents (Amount added)	Thickness of the epidermis (μm)
Bis (diethylene glycol monoethyl ether) fumarate (0.5%)	61.6

(continued)

Agents (Amount added)	Thickness of the epidermis ( $\mu\text{m}$ )
Dimethyl fumarate (0.5%)	40.4
50% ethanol aqueous solution (base)	103.9
No application	127.2

The results of the test show that the addition of fumaric acid diester derivatives inhibited epidermal hyperproliferation as shown in the reduced epidermal thickening by UVB, compared with the group treated with a base alone or the non-application group.

#### Application Example 1 (Skin Cream)

**[0070]** A skin cream was prepared by mixing the fumaric acid diester derivative according to the following composition.

#### (1) Composition

##### **[0071]**

Ingredients	Mixing ratio (mass%)
(Component A)	
• Beeswax	2.0
• Stearic acid	5.0
• Stearyl alcohol	5.0
• Hydrogenated lanolin	2.0
• Squalene	20.0
• Sorbitan monostearate	3.0
• Polyoxyethylene (20) sorbitan monostearate	3.0
• Propylene glycol	5.0
(Component B)	
• Methylparaben	0.2
• Purified water	Balance to the total of 100
(Component C)	
• Bis (diethylene glycol monoethyl ether) fumarate	10.0

#### (2) Preparation method

**[0072]** Bis (diethylene glycol monoethyl ether) fumarate as Component C was added to Component B. After dissolving Components A and B separately by heating them to 80°C, both components were combined and then cooled to 30°C while stirring to make a skin cream.

#### Application Example 2 (Skin Lotion)

**[0073]** A skin lotion was prepared by mixing the fumaric acid diester derivative according to the following composition.

## (1) Composition

**[0074]**

Ingredients	Mixing ratio (mass%)
(Component A)	
· Olive oil	10.0
· Isopropyl myristate	1.0
· Polyoxyethylene (6) nonylphenyl ether	0.5
· Propylene glycol	1.0
· Glycerin	2.0
(Component B)	
· Methylparaben	0.1
· Ethanol	7.0
· Purified water	Balance to the total of 100
(Component C)	
· Bis (diethylene glycol monoethyl ether) fumarate	0.0001

## (2) Preparation method

**[0075]** Bis (diethylene glycol monoethyl ether) fumarate as Component C was added to Component B. After dissolving Components A and B homogeneously, Components A and B were mixed and dispersed with stirring to prepare a lotion. The lotion prepared was poured in a container. The content was shaken to effect a uniform dispersion before use.

## Application Example 3 (Bath Agent)

**[0076]** A bath agent with the following composition was prepared by a common method.

Ingredients	Mixing ratio (mass%)
· Bis (diethylene glycolmonoethyl ether) fumarate	1.0
· Sodium hydrogen carbonate	Balance
· Sodium carbonate	20.0
· Sodium sulfate	15.0
· Sodium chloride	7.5
· Silica	0.5
· 1,3-butylene glycol	1.0
· Urea	1.0
· Seaweed extract	1.0
· Colorant	Appropriate amount
· Dextrin	Appropriate amount
· Perfume	Appropriate amount

## Application Example 4 (Skin Cream)

**[0077]** A skin cream was prepared by mixing the fumaric acid diester derivative and retinoic acid according to the following composition.

## (1) Composition

**[0078]**

Ingredients	Mixing ratio (mass%)
(Component A)	
· Beeswax	2.0
· Stearic acid	5.0
· Stearyl alcohol	5.0
· Hydrogenated lanolin	2.0
· Squalene	20.0
· Sorbitan monostearate	3.0
· Polyoxyethylene (20) sorbitan monostearate	3.0
· Propylene glycol	5.0
(Component B)	
· Methylparaben	0.2
· Purified water	Balance to the total of 100
(Component C)	
· Bis (diethylene glycol monoethyl ether) fumarate	10.0
· Retinoic acid	1.0

## (2) Preparation method

**[0079]** Bis (diethylene glycol monoethyl ether) fumarate and retinoic acid as Component C were added to Component B. After dissolving Components A and B separately by heating to 80°C, both components were combined and then cooled to 30°C with stirring to make a skin cream.

## Application Example 5 (Skin Lotion)

**[0080]** A skin lotion was prepared by mixing the fumaric acid diester derivative and retinoic acid according to the following composition.

## (1) Composition of a skin lotion

**[0081]**

Ingredients	Mixing ratio (mass%)
(Component A)	
· Olive oil	10.0
· Isopropyl myristate	1.0
· Poxoxyethylene (6) nonylphenyl ether	0.5
· Propylene glycol	1.0
· Glycerin	2.0
(Component B)	
· Methylparaben	0.1
· Ethanol	7.0
· Purified water	Balance to the total of 100
(Component C)	
· Bis (diethylene glycol monoethyl ether) fumarate	0.0001
· Retinoic acid	1.0

## (2) Preparation method

**[0082]** Bis (diethylene glycol monoethyl ether) fumarate and retinoic acid as Component C were added to Component B. After dissolving Components A and B homogeneously, Components A and B were mixed and dispersed with stirring to prepare a lotion. The lotion prepared was filled in a container. The contents were shaken to effect uniform dispersion before use.

## Application Example 6 (Skin Lotion)

**[0083]** A skin lotion was prepared by mixing the fumaric acid diester derivative and retinoic acid according to the following composition.

## (1) Composition of a skin lotion

**[0084]**

Ingredients	Mixing ratio (mass%)
(Component A)	
· Olive oil	10.0
· Retinyl acetate	0.02
· Isopropyl myristate	1.0
· Polyoxyethylene (6) nonylphenyl ether	0.5
· Propylene glycol	1.0
· Glycerin	2.0
(Component B)	
· Methylparaben	0.1
· Ethanol	7.0
· Purified water	Balance to the total of 100
(Component C)	
· Bis (diethylene glycol monoethyl ether) fumarate	3.0
· Retinoic acid	1.0

## (2) Preparation method

**[0085]** Bis (diethylene glycol monoethyl ether) fumarate and retinoic acid as Component C were added to Component B. After dissolving Components A and B homogeneously, Components A and B were mixed and dispersed with stirring to prepare a lotion. The lotion prepared was filled in a container. The contents were shaken to effect uniform dispersion before use.

## Application Examples 7 through 9 (Skin Cream)

**[0086]** Skin cream was prepared according to the following composition in a conventional method.

Ingredients	Mixing ratio (mass%)		
	Application Example 7	Application Example 8	Application Example 9
· Bis (diethylene glycol monoethyl ether) fumarate	0.1	0.1	0.1
· Stearic acid	1	1	-
· Isostearic acid	-	-	1
· Glycerin monostearate	2	2	2
· Behenyl alcohol	2	2	2

(continued)

	Ingredients	Mixing ratio (mass%)		
		Application Example 7	Application Example 8	Application Example 9
5	· Bleached beeswax	1	1	-
	· Cetyl myristate	1	1	1
	· Sorbitan sesquioleate	1	1	1
10	· N-stearoylphosphatidylcholine	0.1	0.1	0.1
	· Hydrogenated lecithin	0.1	0.1	0.1
	· Vegetable squalane	5	5	5
15	· Retinyl palmitate	0.05	-	-
	· Octyldodecyl myristate	5	5	5
	· Phellodendron bark extract	0.1	1	0.1
	· <i>Pyracantha fortuneana</i> extract	0.1	0.3	-
20	· Water-soluble glycyrrhiza extract	-	-	0.1
	· 1,3-butylene glycol	5	10	5
	· Conc. Glycerin	5	5	5
25	· Parahydroxybenzoate ester	0.2	0.2	0.2
	· N-acetylglucosamine oligomer	0.1	0.1	0.1
	· Magnesium ascorbylphosphate	0.1	0.1	0.1
	· Sodium ascorbylphosphate	0.1	0.1	0.1
30	· Gamma-aminobutyric acid	0.1	0.1	0.1
	· Sodium N-stearoylglutamate	0.2	0.2	0.2
	· Alkyl-modified carboxyvinyl polymer*1	0.05	0.05	0.05
35	· Nicotinamide	0.1	0.1	0.1
	· Sarcosine	0.1	0.1	0.1
	· Purified water	Balance	Balance	Balance

\*1: PEMULEN™ TR-1, ex BF. Goodrich Company

Application Examples 10 and 11 (Lotion)

[0087] Lotions were prepared according to the following composition in a conventional method.

	Ingredients	Mixing ratio (mass%)	
		Application Example 10	Application Example 11
45	· Bis (diethylene glycol monoethyl ether) fumarate	0.1	0.1
50	· Phellodendron bark extract	0.1	0.3
	· Hibiscus extract	0.2	0.5
	· Lactic acid bacteria fermentation solution	0.1	0.1
55	· 1,3-butylene glycol	5	5
	· Dipropylene glycol	-	5



(continued)

Ingredients	Mixing ratio (mass%)	
	Application Example 10	Application Example 11
· Raffinose	1	1
· Ethanol	-	1
· Phenoxyethanol	0.2	0.2
· Pectin	-	0.05
· Xanthan gum	-	0.1
· Sodium citrate	0.05	0.05
· Horsetail extract	0.1	0.1
· Diisopropylamine dichloroacetate	0.2	0.2
· Gamma-amino-beta-hydroxybutyric acid	0.2	0.2
· Sodium hyaluronate	0.001	0.001
· Dipotassium glycyrrhizinate	0.2	0.2
· Naematoloma sublateralium extract	0.05	0.05
· Decarboxy carnosine hydrochloride	0.05	0.05
· Perfume	0.02	0.02
· Purified water	Balance	Balance

Application Examples 12 through 14 (Gel)

**[0088]** Gels were prepared according to the following composition in a conventional method.

Ingredients	Mixing ratio (mass%)		
	Application Example 12	Application Example 13	Application Example 14
· Bis (diethylene glycol monoethyl ether) fumarate	0.1	0.1	0.1
· Decamethylcyclopentasiloxane	10	10	10
· Isostearyl isostearate	1	-	-
· Olive oil	-	1	-
· Macadamianut oil	-	-	1
· Eucalyptus oil	0.1	-	0.1
· Hexyldecanol	1	0.1	-
· dl-alpha-tocopheryl nicotinate	-	0.1	-
· Retinol	0.1	-	-
· Polyoxyethylene (60) hydrogenated castor oil	2	2	2
· Spherical silicone powder*2	1	1	5
· Phellodendron bark extract	0.1	1	0.1
· Water-soluble chlorophyll	0.02	0.02	0.02
· Salvia extract	-	0.3	0.1

\*2 Tospearl™, ex GE Toshiba Silicone Corp.

(continued)

Ingredients	Mixing ratio (mass%)		
	Application Example 12	Application Example 13	Application Example 14
· 1,3-butylene glycol	5	10	5
· Sorbitol solution	3	3	3
· Polyethylene glycol 4000	1	1	1
· Carboxyvinyl polymer	0.2	0.2	0.2
· Glycoceramide* <sup>3</sup>	0.1	0.1	0.1
· Parahydroxybenzoate ester	0.2	0.2	0.2
· Mevalonolactone	0.5	0.5	0.5
· Edetic acid salt	0.02	0.02	0.02
· Potassium hydroxide	0.05	0.05	0.05
· Purified water	Balance	Balance	Balance

\*<sup>3</sup> Bioceramide™, ex Kibun Food Chemical Corp.

Application Examples 15 through 17 (Lipophilic Cream)

[0089] Lipophilic creams were prepared according to the following composition in a conventional method.

Ingredients	Mixing ratio (mass %)		
	Application Example 15	Application Example 16	Application Example 17
· Bis (diethylene glycol monoethyl ether) fumarate	0.1	0.1	0.1
· Comodified silicone* <sup>4</sup>	2	2	2
· Polyoxyethylene modified silicone dispersion* <sup>5</sup>	-	2	-
· Squalane	-	-	10
· Decamethylcyclopentasiloxane	15	20	10
· Retinol	0.05	0.05	0.05
· Methylpolysiloxane	5	2	3
· Cholesteryl long-chain branched fatty acids ester* <sup>6</sup>	-	-	3
· Silicone elastomer dispersion* <sup>7</sup>	5	2	-
· Phellodendron bark extract	1	1	1
· Glycyrrhiza extract	0.1	0.1	0.1
· Water-soluble chlorophyll	0.02	0.02	0.02
· Sodium chloride	1	1	1
· Dipropylene glycol	5	5	5
· Conc. Glycerin	5	5	5

\*<sup>4</sup>: ABIL™ EM 90, ex Goldschmidt AG.\*<sup>5</sup>: Silicone BY22-008, ex Toray Dow Corning Silicone Co.\*<sup>6</sup>: YOFCO CLE-NH, ex Nippon Seika Corporation Co.\*<sup>7</sup>: Trefil™, ex Toray Dow Corning Silicone Co.

(continued)

Ingredients	Mixing ratio (mass %)		
	Application Example 15	Application Example 16	Application Example 17
· Raffinose	1	1	1
· Parahydroxybenzoate ester	0.3	0.3	0.3
· N-methyl-L-serine	0.5	0.5	0.5
· Purified water	Balance	Balance	Balance

Application Examples 18 through 20 (Sunscreen)

[0090] Sunscreens were prepared according to the following composition in a conventional method.

Ingredients	Mixing ratio (mass%)		
	Application Example 19	Application Example 20	Application Example 21
· Bis (diethylene glycol monoethyl ether) fumarate	0.1	0.1	0.1
· Dioctyl ether	22	15	10
· Comodified silicone* <sup>4</sup>	2	2	2
· Glyceryl tri-2-ethylhexanoate	-	-	5
· Hydrogenated oil	-	-	0.1
· Methylphenyl polysiloxane	-	3	-
· Macadamianut fatty acid phytosteryl ester	-	-	2
· 2-ethylhexyl 4-methoxycinnamate	-	7	7
· Titanium oxide	5	-	4
· Zinc oxide	5	-	4
· Phellodendron bark extract	1	1	1
· Magnesium chloride	1	1	1
· 1,3-butylene glycol	5	5	5
· Phenoxyethanol	0.3	0.3	0.3
· Hibiscus extract	1	1	1
· Aloe extract	0.1	0.1	0.1
· Yeast extract* <sup>8</sup>	1	1	1
· Purified water	Balance	Balance	Balance

\*<sup>4</sup>: ABIL™ EM90, ex Goldschmidt AG.\*<sup>8</sup>: DISMUTIN™, ex Pentapharm AG.


## Industrial Applicability

[0091] The present invention provides an epidermal barrier function-enhancing agent and a composition for skin external application which are very safe, quickly ameliorate the epidermal permeability barrier function disruption, and have excellent effects on maintaining the skin in a cosmetologically healthy state; a HB-EGF expression-inducing agent to enhance epidermal turnover; an amphiregulin expression inhibitor to diminish epidermal hyperproliferation; an agent to be used together with retinoic acid or a derivative thereof to prevent dry skin caused by retinoic acid or a derivative

thereof; and an ameliorating agent for aged skin which ameliorates metabolic disruption and skin thinning of the skin due to aging.


## Claims

1. A skin external agent composition, **characterized in that** it comprises a fumaric acid diester derivative expressed by the following general formula (1)




, wherein R<sup>1</sup> represents a C<sub>1-4</sub> alkylene group; R<sup>2</sup> and R<sup>3</sup> each represents a linear or branched C<sub>1-8</sub> alkyl group; and n is an integer of 2 to 5.

2. The Skin external agent composition as described in Claim 1, wherein the composition is a cosmetic.
3. An epidermal permeability barrier function-enhancing agent comprising a fumaric acid diester derivative expressed by the following general formula (1)



, wherein R<sup>1</sup> represents a C<sub>1-4</sub> alkylene group; R<sup>2</sup> and R<sup>3</sup> each represents a linear or branched C<sub>1-8</sub> alkyl group; and n is an integer of 2 to 5.


4. An amphiregulin expression inhibitor comprising a fumaric acid diester derivative expressed by the following general formula (1)



, wherein R<sup>1</sup> represents a C<sub>1-4</sub> alkylene group; R<sup>2</sup> and R<sup>3</sup> each represents a linear or branched C<sub>1-8</sub> alkyl group;


and n is an integer of 2 to 5.

5. An epidermal excessive proliferation inhibitor comprising a fumaric acid diester derivative expressed by the following general formula (1)




, wherein R<sup>1</sup> represents a C<sub>1-4</sub> alkylene group; R<sup>2</sup> and R<sup>3</sup> each represents a linear or branched C<sub>1-8</sub> alkyl group; and n is an integer of 2 to 5.

6. A heparin-binding epidermal growth factor-like growth factor (HB-EGF) expression-inducing agent comprising a fumaric acid diester derivative expressed by the following general formula (1)




, wherein R<sup>1</sup> represents a C<sub>1-4</sub> alkylene group; R<sup>2</sup> and R<sup>3</sup> each represents a linear or branched C<sub>1-8</sub> alkyl group; and n is an integer of 2 to 5.

7. An ameliorating agent for aged skin comprising a fumaric acid diester derivative expressed by the following general formula (1)




, wherein R<sup>1</sup> represents a C<sub>1-4</sub> alkylene group; R<sup>2</sup> and R<sup>3</sup> each represents a linear or branched C<sub>1-8</sub> alkyl group; and n is an integer of 2 to 5.

8. An agent to be used together with retinoic acid or a derivative thereof, said agent comprising a fumaric acid diester derivative expressed by the following general formula (1)




, wherein R<sup>1</sup> represents a C<sub>1-4</sub> alkylene group; R<sup>2</sup> and R<sup>3</sup> each represents a linear or branched C<sub>1-8</sub> alkyl group; and n is an integer of 2 to 5.

9. An agent to prevent dry skin caused by retinoic acid or a derivative thereof, said agent comprising a fumaric acid diester derivative expressed by the following general formula (1)



, wherein R<sup>1</sup> represents a C<sub>1-4</sub> alkylene group; R<sup>2</sup> and R<sup>3</sup> each represents a linear or branched C<sub>1-8</sub> alkyl group; and n is an integer of 2 to 5.

10. An epidermal external agent composition comprising a fumaric acid diester derivative expressed by the following general formula (1)



, wherein R<sup>1</sup> represents a C<sub>1-4</sub> alkylene group; R<sup>2</sup> and R<sup>3</sup> each represents a linear or branched C<sub>1-8</sub> alkyl groups; and n is an integer of 2 to 5, and retinoic acid or a derivative thereof.